


Relevant Antiproliferative Activity against *Toxoplasma gondii* of a new binuclear Co(II) complex

Vagner M. de Assis^{1,2*}, Luana C. Batista², Lorenzo C. Visentin³, Fernanda S. de Souza⁴, Renato A. DaMatta⁴, Adolfo Horn², Christiane Fernandes²

¹Instituto Federal Fluminense, Campus Cabo Frio, RJ – Brasil, ²Laboratório de Ciências Químicas, Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, RJ – Brasil, ³P&D NanoBusiness Informação e Inovação, Duque de Caxias, RJ, Brazil, ⁴Laboratório de Biologia Celular e Tecidual, Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, RJ, Brazil

*e-mail: vagnerassis@gmail.com

Cobalt is an essential element for life and it is less toxic than non-essential metals such as platinum. Cobalt is an integral component of cyanocobalamin (vitamin B12)¹. Several recent publications in the literature highlight the relevant properties of cobalt complexes as antitumor compounds². Other studies report the antibacterial and antiviral activities of cobalt complexes³. Recently, we report the cytotoxic effect on *Toxoplasma gondii* of a dinuclear iron(III) complex [Fe(HPCINOL)(SO₄)₂]-μ-oxo, which induces reduction of superoxide dismutase and catalase activities disturbing the parasite redox equilibrium, resulting in cystogenesis and parasite death⁴. In this work we report the synthesis, characterization and antiproliferative activity against an intracellular parasite of a new binuclear cobalt(II) complex, containing the ligand HPCINOL= 1-(bis-pyridin-2-ylmethyl-amino)-3-chloropropan-2-ol, which was previously described⁵. The binuclear complex [(HPCINOL)Co(-μ-Cl₂)Co(HPCINOL)](ClO₄)₂ **1** which was characterized by a variety of physic-chemical methods. M.p.: 230 °C. Anal. Calc. for C₃₀H₃₆Cl₆Co₂N₆O₁₀; MW = 971 g mol⁻¹: C, 37.10; H, 3.74; N, 8.85. Found: C, 37.47; H, 3.58; N, 8.85. Other physic-chemical methods were used, such as electronic spectroscopy, ESI(+)-MS, ESI(+)-MS/MS and X-ray diffraction, which allowed the resolution of complex structure (Fig. 1), which is a binuclear cation containing two counter-ions perchlorate and has octahedral structure distorted in their metal centers. Complex **1** was non-toxic to the host cells. This complex presented high anti-toxoplasma effect in infected LLC-MK₂ host cells. Comparing



with control (no treatment), the ligand and Sulfadiazine did not control parasite growth after 24 h of treatment. However, the complex reduced parasite growth by 55% when 10 μmol L⁻¹ was used. After 48 h of treatment this complex presented a reduction of *T. gondii* growth similar to 24 h.

Figure 1. ORTEP for complex **1**.

- 1- Andreini, C.; Bertini, I.; Cavallaro, G.; Holliday, G.L.; Thornton, J.M. J. Biol. Inorg. Chem. 2008, 13, 1205–1218.
- 2- Munteanu, C.R.; Suntheralingam, K. Dalton Trans., 2015, 44 (31), 13796–13808.
- 3- Chang, E.L.; Simmers, C.; Knight, D.A. Pharmaceuticals, 2010, 3, 1711–1728.
- 4- Portes, J.A.; Souza, T.G.; dos Santos, T.A.T.; Silva, L.L.R.; Ribeiro, T.P.; Pereira, M.D.; Horn Jr., A.; Fernandes, C.; DaMatta, R.A.; Souza, W.; Seabra, S.H. Antimicrob. Agents Chemother. 2015, 59, 7374–7386.
- 5- Horn Jr., A.; Fernandes, C.; Bortoluzzi, A.J.; Vugman, N.V.; Herbst, M.H. J. Mol. Struct. 2005, 749, 96–102.