

# Zinc Fingers as Templates for Platinum-Metal Coordination Chemistry

S.D. Tsotsoros<sup>1</sup>, R. E.F. da Paiva<sup>2</sup>, Z.Du<sup>1</sup>, E. J. Peterson<sup>1</sup>, C. Abbehausen<sup>2</sup>, P.P. Corbi<sup>2</sup>, V.H.F. Bernardes<sup>3</sup>, F.A. Lima<sup>2</sup>, M.D. Vargas<sup>3</sup> and N.P. Farrell<sup>1,\*</sup>


<sup>1</sup>Department of Chemistry, Virginia Commonwealth University, Richmond, USA

<sup>2</sup>Department of Chemistry, University of Campinas, Campinas SP, Brazil and

<sup>3</sup>Department of Chemistry, Federal Fluminense University, Niteroi RJ, Brazil.

Email: [npfarrell@vcu.edu](mailto:npfarrell@vcu.edu)

This lecture summarises our work on the fundamental chemistry and biology of platinum-metal (Pt, Pd and Au) complexes with zinc finger proteins. Zinc finger proteins are involved in a host of biological processes such as transcription and DNA repair and they are important targets for therapeutic intervention in viral diseases and cancer. A specific focus will be on the HIV NCp7 nucleocapsid protein (NC), a small basic nucleic acid chaperone containing two zinc finger (or zinc knuckle) Cys<sub>3</sub>His motifs and its interaction with RNA. Conceptual analogies between alkylation and metallation in biology have been especially useful in suggesting new patterns of platinum-metal and platinum-nucleobase chemistry. These patterns include molecular recognition and electrophilic attack by platinum-metal compounds on zinc-thiolate bonds. Protonation, alkylation or coordination to a metal ion, such as Pt(II) or Au(III) of a nucleobase generally decreases the energy of its lowest unoccupied molecular orbital (LUMO), thereby improving the potential for  $\pi$ -stacking interactions with the highest occupied molecular orbital (HOMO) in aromatic aminoacids such as N-acetyl tryptophan [1,2]. This effect has been used to target platinum-metal-nucleobases to the tryptophan-containing “full” zinc finger and its single C-terminal peptide. Fluorescence quenching assay, NMR and Circular Dichroism (CD) spectroscopy, ESI-Mass Spectrometry (ESI-MS) and gel-shift electrophoresis show that these small molecules may be effective inhibitors of the nucleocapsid-nucleic acid interaction. X-Ray Absorption Spectroscopy studies can distinguish between Au(I) and Au(III) binding to the peptide. The factors affecting selective “non-covalent” ligand-biomolecule recognition through enhanced  $\pi$ - $\pi$  stacking between metallated nucleobases and the tryptophan-containing nucleocapsid will be examined. In a second step, such recognition can lead to formation of the “covalent” metal-thiolate bond and subsequent zinc ejection. These findings and others suggest that zinc finger proteins are excellent templates for metal ion exchange and ligand reactivity [3].



1. Anzellotti, A., Liu, Q., Bloemink, M.J., Scarsdale, J.N. and Farrell, N. (2006) Targeting the Retroviral Zinc Finger-DNA Interaction. A small molecule approach utilizing the electrophilic nature of trans-Platinum-Nucleobase Compounds. *Chemistry and Biology*. 13:539-548
2. Anzellotti, A.I., Bayse, C.A. and Farrell, N.P. (2008) Effects of Nucleobase Metallation on Frontier Molecular Orbitals: *Inorg. Chem.* 47:10425-10431.
3. Quintal, S.M., dePaula, Q.A. and Farrell, N.P. (2011) Zinc Finger Proteins as Templates for Metal Ion Exchange and Ligand Reactivity. Chemical and Biological Consequences. *Metallomics*. 3:121-139.

This work was supported by National Science Foundation NSF CHE-1413189, São Paulo Research Foundation (FAPESP) 2014/20334-0 and 2015/9905-1, Brazilian Federal Agency for the Support and Evaluation of Graduate Education (CAPES), CAPES/PVES 154/2012 and 0580/2013. FAL acknowledges National Council for Scientific and Technological Development (CNPq) for the productivity grant 311270/2015-8.