

Reaction of Au(I) complexes with HIV-1 and human zinc finger proteins

X-ray absorption spectroscopy

Saulo Q. Gomes^{1*}, Raphael E. F. de Paiva¹, Frederico A. Lima², Zhifeng Du³, Camilla Abbehausen¹, Pedro P. Corbi¹, Nicolas P. Farrell³

¹*Institute of Chemistry, University of Campinas – UNICAMP, Campinas, São Paulo, Brazil,*

²*Centro Nacional de Pesquisa em Energia e Materiais, Brazilian Synchrotron Light Laboratory – LNLS, Campinas, SP, Brazil,* ³*Department of Chemistry, Virginia Commonwealth University, Richmond, Virginia, United States*

*e-mail: saulo.gomes@iqm.unicamp.br

Current antiretroviral therapies for HIV-1 infection can be considered efficient because they increase life expectancy, but on the other hand, are ineffective to eradicate the infection and resistant strains emerge.¹ To overcome these problems new molecular targets are investigated and new inhibitors are needed to maintain the therapies functional. A recognized molecular target for HIV-1 therapy is the zinc finger (ZnF) protein 7 of the nucleocapsid (NCp7).² Gold complexes have been showing inhibition of NCp7 by electrophilic attack of the metal to cysteines leading to the ejection of Zn(II) and formation of so called gold fingers.^{3,4} However, structural details of the final products are currently lacking. Early studies on anti-arthritis gold(I)-based drug, auranofin, showed to decrease the viral load of HIV-1 in an infected patient.¹ and recently in studies performed in SIV infected rhesus macaques.¹ Recently our research group studied the interaction of [Ph₃PAuL] complexes with NCp7 finding the first formation of Ph₃PAu-apo-peptide as a stable species.⁴

X-ray absorption spectroscopy studies have allowed the development of a model for the coordination environment of gold fingers based on the analysis of gold L₃-edge XAS of ZnF+Au systems, and simulations of XANES spectra. The final products of [AuCl(PEt₃)] and auranofin with the C-terminal finger of NCp7 can be definitively categorized as containing S-Au-P coordination spheres, confirming the Lewis acid electrophilic attack on the zinc-bound cysteines. This conclusion was confirmed by *ab initio* multiple scattering simulations of the XANES spectra of hypothetical compounds with Et₃P-Au-His, Et₃P-Au-Cys, Et₃P-Au-PEt₃ and [AuCl(PEt₃)] itself. XAS could also be used to distinguish between limiting mechanisms involving structurally distinct zinc fingers – the reaction product of [AuCl(PEt₃)] with Sp1 is different from that formed in the reaction of the former with NCp7, containing only gold finger. The XAS spectrum indicating the presence of a more electron rich gold center and a S-Au-S coordination environment for the final product of this reaction was proposed. This result corroborates to the design of more selective compounds.

1. Lewis, M. G.; Dafonseca, S.; Chomont, N.; Palamara, A. T.; Tardugno, M.; Mai, A.; Collins, M.; Wagner, W. L.; Yalley-ogunro, J.; Greenhouse, J.; Chirullo, B.; Norelli, S.; Garaci, E.; Savarino, A. ;*AIDS* **2011**, 25, 1347.
2. de Rocquigny, H.; Shvadchak, V.; Avilov, S.; Dong, C. Z.; Dietrich, U.; Darlix, J.-L.; Mély, Y. ;*Mini Rev. Med. Chem.* **2008**, 8, 24.
3. Larabee, J. L.; Hocker, J. R.; Hanas, J. S. ;*Chem. Res. Toxicol.* **2005**, 18, 1943.
4. Abbehausen, C.; Peterson, E. J.; De Paiva, R. E. F.; Corbi, P. P.; Formiga, A. L. B.; Qu, Y.; Farrell, N. P. ;*Inorg. Chem.* **2013**, 52, 11280.