

# Synthesis, structural characterization, high resolution mass spectrometric studies and antimicrobial assays of a silver complex with sulfisoxazole

Ana Thereza M. Fiori<sup>1\*</sup>, Alexandre Cuin<sup>2</sup>, Wilton R. Lustri<sup>3</sup>, Pedro P. Corbi<sup>1</sup>

<sup>1</sup> *Institute of Chemistry, University of Campinas – UNICAMP, Campinas-SP Brazil,*

<sup>2</sup> *Department of Chemistry, Exact Sciences Institute – UFJF, Juiz de Fora-MG Brazil,*

<sup>3</sup> *Biological and Health Sciences Department – UNIARA, Araraquara-SP, Brazil*

\*e-mail: anatherezaefiori@gmail.com

The discovery of the antibacterial activity of sulfonamides and their applications in the treatment of various infections changed the antimicrobial therapy that we know today, bringing a significant improvement in quality of life.<sup>1,2</sup> However, the excessive and indiscriminate use of sulfa drugs and other classes of antimicrobial agents led to the emergence of antibiotic-resistant bacteria, turning the search for new drugs extremely important.<sup>1-4</sup> One of the strategies to develop new antibacterial agents, is the coordination of drugs with metals, both showing recognized antibacterial activities.<sup>1,2,4</sup> In this context, the present work describes the synthesis, structural characterization and biological assays of a new silver(I) complex with the sulfa drug sulfisoxazole (SIZ). Elemental, thermal and high resolution mass spectrometric studies (ESI-QTOF-MS) led to a 1:1 metal/ligand composition, suggesting the minimal formula  $\text{AgC}_{11}\text{H}_{12}\text{O}_3\text{N}_3\text{S}\cdot 0.5\text{H}_2\text{O}$  (Ag-SIZ). Infrared and solution state NMR spectroscopic measurements indicated ligand coordination to Ag(I) by the nitrogen atoms of the  $\text{SO}_2\text{N}$  and  $\text{NH}_2$  groups. This hypothesis was confirmed by crystallographic studies, which shows the formation of a dimer with two Ag(I) bridging between two SIZ molecules. Antibacterial assays indicated that the Ag-SIZ complex is effective against Gram-negative (*E. coli* and *P. aeruginosa*) and Gram-positive (*S. aureus*). The complex was shown to be more active against the Gram-positive bacterial strain.

<sup>1</sup> Penildon, S. In *Sulfonamidas e outros quimioterápicos empregados no tratamento de infecções do trato urinário*. Carvalho, W. A.; Guanabara Koogan: Rio de Janeiro, Brazil, 2006, ch. 102.

<sup>2</sup> Rocha, D. P.; Pinto, G. F.; Ruggiero, R.; Oliveira, C. A.; Guerra, W.; Fontes, A. P. S.; Tavares, T. T.; Marzano, I. M.; Maia, E. C. P. *Quím. Nova*. 2011, 34, 111.

<sup>3</sup> Zanini, A. C.; Oga S. In *Sulfonamidas antimicrobianas*. Silva, A. C. G.; Atheneu: São Paulo, Brazil, 1985, ch. XXXIII.

<sup>4</sup> Nazari, Z. E.; Banoee, M.; Sepahi, A. A.; Rafii, F.; Shahverdi, A. R. *Gold Bull.* 2012, 45, 53.

<sup>5</sup> Silva, A. C. G. *Farmacologia Aplicada*, 3<sup>th</sup> ed.; Atheneu: São Paulo, Brazil, 1985.

This study was supported by grants from the Brazilian Agencies FAPESP (grant#2015/25114-4), CNPq (grant#442123/2014-0) and FAPEMIG.